

Bicomplexes and finite Toda lattices

A. Dimakis¹ and F. Müller-Hoissen²

¹ Department of Mathematics, University of the Aegean
GR-83200 Karlovasi, Samos, Greece

² Max-Planck-Institut für Strömungsforschung
Bunsenstrasse 10, D-37073 Göttingen, Germany

Abstract

We associate bicomplexes with the finite Toda lattice and with a finite Toda field theory in such a way that conserved currents and charges are obtained by a simple iterative construction.

In recent work [1, 2] we have demonstrated how bicomplexes can be associated with several completely integrable models in such a way that they provide us with an iterative construction of conserved currents. In the following we recall the underlying mathematical structure and demonstrate how the finite Toda chain and a finite Toda field equation fit into this scheme.

Let $V = \bigoplus_{r \geq 0} V^r$ be an $\mathbb{N}_0$ -graded linear space (over $\mathbb{R}$ or $\mathbb{C}$) and $d, \delta : V^r \rightarrow V^{r+1}$ two linear maps satisfying $d^2 = 0$, $\delta^2 = 0$ and $d\delta + \delta d = 0$. Then (V, d, δ) is called a *bicomplex*. In the following we assume that, for some $s \in \mathbb{N}$, $H_\delta^s(V)$ is trivial, so that all δ -closed elements of V^s are δ -exact. Furthermore, we assume that there is a (nonvanishing) $\chi^{(0)} \in V^{s-1}$ with $dJ^{(0)} = 0$ where $J^{(0)} = \delta\chi^{(0)}$. Let us define $J^{(1)} = d\chi^{(0)}$. Then $\delta J^{(1)} = -d\delta\chi^{(0)} = 0$, so that $J^{(1)} = \delta\chi^{(1)}$ with some $\chi^{(1)} \in V^{s-1}$. Next we define $J^{(2)} = d\chi^{(1)}$. Then $\delta J^{(2)} = -d\delta\chi^{(1)} = -dJ^{(1)} = -d^2\chi^{(0)} = 0$, so that $J^{(2)} = \delta\chi^{(2)}$ with some $\chi^{(2)} \in V^{s-1}$. This can be iterated further and leads to a (possibly infinite) chain (see Fig. 1) of δ -closed elements $J^{(m)}$ of V^s and $\chi^{(m)} \in V^{s-1}$ satisfying

$$J^{(m+1)} = d\chi^{(m)} = \delta\chi^{(m+1)}. \quad (1)$$

Introducing $\chi = \sum_{m \geq 0} \lambda^m \chi^{(m)}$ with a parameter λ , this becomes

$$\delta(\chi - \chi^{(0)}) = \lambda d\chi. \quad (2)$$

In the following examples we will only consider the case where $s = 1$. In these examples, the $J^{(m)}$ represent conserved currents which determine conserved charges.




Fig. 1

The chain of δ -closed s -forms $J^{(m)}$.

Example 1: The finite Toda lattice.

Let $C^\infty(\mathbb{R}, \mathbb{R}^n)$ be the set of smooth maps $f : \mathbb{R} \rightarrow \mathbb{R}^n$ and $\Lambda = \bigoplus_{r=0}^2 \Lambda^r$ the exterior algebra of a 2-dimensional vector space. We choose linearly independent 1-forms $\tau, \xi \in \Lambda^1$ which satisfy $\tau^2 = \xi^2 = \tau\xi + \xi\tau = 0$. Furthermore, we set $V = \bigoplus_{r=0}^2 V^r = C^\infty(\mathbb{R}, \mathbb{R}^n) \otimes \Lambda$. Next we define linear maps $d, \delta : V^0 \rightarrow V^1$ via

$$df = (Lf)\tau + (\dot{f} + Mf)\xi, \quad \delta f = \dot{f}\tau + (Sf - f)\xi \quad (3)$$

with maps $S, M, L : \mathbb{R} \rightarrow M(n \times n; \mathbb{R})$ and $\dot{f} = df/dt$ where t denotes the canonical coordinate function on $\mathbb{R}$. d extends to V^1 via $d(f\tau + g\xi) = (df)\tau + (dg)\xi$ for all $f, g \in C^\infty(\mathbb{R}, \mathbb{R}^n)$, and correspondingly for δ . d and δ satisfy the bicomplex conditions iff

$$\dot{S} = 0, \quad \dot{M} = [S, L], \quad \dot{L} = [L, M]. \quad (4)$$

Let $e_i = (\delta_i^a)$ denote the standard basis of $\mathbb{R}^n$ and $E_{ij} = (\delta_i^a \delta_{jb})$ the elementary matrices. Then we have $E_{ij}e_k = \delta_{jk}e_i$ and $E_{ij}E_{kl} = \delta_{jk}E_{il}$. Now we choose

$$S = \sum_{i=1}^{n-1} E_{i,i+1}, \quad M = \sum_{i=1}^n \dot{q}_i E_{ii}, \quad L = - \sum_{i=1}^{n-1} e^{q_i - q_{i+1}} E_{i+1,i}. \quad (5)$$

S has the properties $\dot{S} = 0$, $Se_1 = 0$, $Se_i = e_{i-1}$ for $i = 2, \dots, n$. One finds that $\dot{L} = [L, M]$ is identically satisfied and $\dot{M} = [S, L]$ is equivalent to the finite Toda lattice equation

$$\ddot{q}_1 = -e^{q_1 - q_2}, \quad \ddot{q}_n = e^{q_{n-1} - q_n}, \quad \ddot{q}_i = e^{q_{i-1} - q_i} - e^{q_i - q_{i+1}} \quad i = 2, \dots, n-1. \quad (6)$$

$\delta J = 0$ for $J = J_0\tau + J_1\xi$ means $SJ_0 - J_0 = \dot{J}_1$. In particular, this implies $J = \delta\phi$ with $\phi = \sum_{k=1}^n \phi_k e_k$ and $\phi_k = -\sum_{i=k}^n J_{1i}$. Hence δ -closed elements of V^1 are δ -exact. Using the euclidean scalar product $\langle \cdot, \cdot \rangle$ and $u = \sum_{i=1}^n e_i$, we define

$$Q = \langle u, J_1 \rangle \quad (7)$$

for a δ -closed element $J \in V^1$. Then

$$\dot{Q} = \langle u, SJ_0 - J_0 \rangle = \langle S^t u - u, J_0 \rangle = -\langle e_1, J_0 \rangle. \quad (8)$$

The δ -closed elements $J^{(m)} \in V^1$ obtained via the above iteration procedure (for $s = 1$) satisfy $J^{(m)} = d\chi^{(m-1)}$ which implies $J_0^{(m)} = L\chi^{(m-1)}$ and thus $\dot{Q}^{(m)} = 0$ since $L^t e_1 = 0$. Hence, the $Q^{(m)}$ are conserved.

Choosing $\chi^{(0)} = u$ (which satisfies $d\delta\chi^{(0)} = 0$), the linear equation (2) becomes equivalent to the system

$$\dot{\chi} = \lambda L\chi, \quad (I - S)\chi = e_n - \lambda(\lambda L\chi + M\chi) \quad (9)$$

where I denotes the $n \times n$ unit matrix. Using $(I - S)^{-1}e_k = \sum_{j=1}^k e_j$, the last equation allows the recursive calculation of the $\chi^{(m)}$:

$$\chi^{(1)} = -(I - S)^{-1}M\chi^{(0)} = -\sum_{i=1}^n \sum_{j=1}^i \dot{q}_i e_j = -\sum_{j=1}^n \sum_{i=j}^n \dot{q}_i e_j \quad (10)$$

$$\chi^{(m)} = -(I - S)^{-1}(M\chi^{(m-1)} + L\chi^{(m-2)}) \quad m = 2, \dots, n. \quad (11)$$

From $J^{(m)} = d\chi^{(m-1)}$ we obtain in particular

$$J^{(1)} = -\sum_{i=2}^n e^{q_{i-1}-q_i} e_i \tau + \sum_{i=1}^n \dot{q}_i e_i \xi \quad (12)$$

$$J^{(2)} = -\sum_{k=1}^{n-1} e^{q_k-q_{k+1}} \chi_k^{(1)} e_{k+1} \tau + \sum_{k=1}^n (\dot{\chi}_k^{(1)} + \dot{q}_k \chi_k^{(1)}) e_k \xi. \quad (13)$$

The associated conserved charges are

$$Q^{(1)} = -\sum_{k=1}^n \dot{q}_k, \quad Q^{(2)} = -\frac{1}{2} \sum_{k=1}^n \dot{q}_k^2 - \sum_{k=1}^{n-1} e^{q_k-q_{k+1}} - \frac{1}{2} (Q^{(1)})^2. \quad (14)$$

To obtain the last expression, we made use of the equations of motion (6). The conserved charges of the finite Toda lattices are well-known, of course [3]. On the n -point lattice, there are only n independent conserved charges. ■

Example 2: Finite Toda field theory.

Modifying the previous example, we now consider $V = C^\infty(\mathbb{R}^2, \mathbb{R}^n) \otimes \Lambda$ and define linear maps $d, \delta : V^r \rightarrow V^{r+1}$, $r = 0, 1$, via

$$df = (Lf)\tau + (Mf - f_x)\xi, \quad \delta f = f_t \tau + (Sf - f)\xi \quad (15)$$

in terms of coordinates t, x on $\mathbb{R}^2$. f_t and f_x denote the partial derivative with respect to t and x , respectively. The bicomplex conditions are then equivalent to

$$S_t = 0, \quad M_t = [S, L], \quad L_x = -[L, M]. \quad (16)$$

Choosing

$$S = \sum_{i=1}^{n-1} E_{i,i+1}, \quad M = - \sum_{i=1}^n q_{i,x} E_{ii}, \quad L = - \sum_{i=1}^{n-1} e^{q_i - q_{i+1}} E_{i+1,i}, \quad (17)$$

we obtain the field equations

$$\begin{aligned} q_{1,tx} &= e^{q_1 - q_2}, & q_{n,tx} &= -e^{q_{n-1} - q_n}, \\ q_{i,tx} &= e^{q_i - q_{i+1}} - e^{q_{i-1} - q_i} & i &= 2, \dots, n-1. \end{aligned} \quad (18)$$

In the following we restrict our consideration to the case $n = 2$ where this system is equivalent to the Liouville equation. Indeed, setting $q_1 = -q_2 = \phi$, we have

$$S = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M = \phi_x \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad L = -e^{2\phi} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (19)$$

and the field equations reduce to

$$\phi_{tx} = e^{2\phi}. \quad (20)$$

Writing $\chi = (a, b)^t$, the linear system (2) with $\chi^{(0)} = u$ yields

$$a_t = 0, \quad b_t = -\lambda a e^{2\phi} \quad (21)$$

and

$$b = 1 + \lambda (b_x - \phi_x b), \quad b = a - \lambda (a_x + \phi_x a). \quad (22)$$

As a consequence, we have

$$a - 1 - 2\lambda a_x + \lambda^2 (a_{xx} + v a) = 0 \quad (23)$$

where we introduced

$$v = \phi_{xx} - \phi_x^2 \quad (24)$$

which satisfies $v_t = 0$ as a consequence of the Liouville equation.¹ The conserved current is

$$J = d\chi = -e^{2\phi} a \begin{pmatrix} 0 \\ 1 \end{pmatrix} \tau - \begin{pmatrix} a_x + \phi_x a \\ b_x - \phi_x b \end{pmatrix} \xi \quad (25)$$

and the associated conserved charge (constructed as in the previous example) becomes

$$Q = -(a_x + \phi_x a) - (b_x - \phi_x b) = \lambda^{-1} (1 - a) \quad (26)$$

¹Actually $v_t = 0$ is equivalent to the Liouville equation as long as $\phi_{tx} \neq 0$. Regarding (24) as an equation for $w = \phi_x$ with a given function $v(x)$, we recover the Ricatti equation $w_x - w^2 - v(x) = 0$ associated with the Liouville equation.

with the help of (22). Note that $a_t = 0$ implies directly $Q_t = 0$. The corresponding conserved charges $Q^{(m)}$ are

$$Q^{(1)} = 0, \quad Q^{(2)} = v, \quad Q^{(3)} = 2v_x, \quad Q^{(4)} = 3v_{xx} - v^2 \quad (27)$$

and so forth. From the above formula for a it is obvious that all the $Q^{(m)}$ depend on ϕ only through v . Since v is conserved, the $Q^{(m)}$ are trivially conserved in this example.

Above we used a special solution of $d\delta\chi^{(0)} = 0$. Its general solution is

$$a^{(0)} = c_1(x) + \int^t c_2(t') e^{-\phi} dt' \quad (28)$$

$$b^{(0)} = \left(c_4(t) + \int^x [c_3(x') - \int^t a^{(0)} e^{2\phi} dt'] e^{-\phi} dx' \right) e^{\phi} \quad (29)$$

with arbitrary differentiable functions $c_1, \dots, c_4$ of a single variable. (26) has to be replaced by $Q = \lambda^{-1}(a^{(0)} - a)$. Our construction of conserved quantities does not really lead to anything new, however. In particular, we get $Q^{(1)} = h(x)$ with an arbitrary function of x (related to c_3) and $Q^{(2)} = c_{1,xx} + 2h_x + c_1 v$. The corresponding recursion formula is $Q^{(m)} = 2Q_x^{(m-1)} - Q_{xx}^{(m-2)} - vQ^{(m-2)}$ for $m > 1$. ■

We expect that convenient bicomplexes can also be associated with generalizations (see [4], for example, and the references therein) of the above Toda systems in order to generate their conserved currents and charges.

Acknowledgment

F. M.-H. profitted quite a lot from participation in a long series of conferences on modern topics in mathematical physics organized by Professor H.-D. Doebner and his collaborators over the years. On the occasion of Professor Doebner's retirement, it is a pleasure to thank him for his support and his outstanding promotion of mathematical physics especially in Germany.

References

- [1] A. Dimakis and F. Müller-Hoissen, Bi-differential calculi and integrable models, math-ph/9908015.
- [2] A. Dimakis and F. Müller-Hoissen, Bi-differential calculus and the KdV equation, math-ph/9908016, to appear in *Rep. Math. Phys.*.
- [3] M. Hénon, Integrals of the Toda lattice, *Phys. Rev. B* **9** (1974) 1921-1923.
- [4] L. Chao, Toda-like systems: solutions and symmetries, hep-th/9512198.